arXiv:nlin/0509027v1 [nlin.SI] 12 Sep 2005

LAX REPRESENTATION OF WZNW-LIKE SYSTEMS

A.V. Balandin [1]
Department of Mathematics and Mechanics
N.I.Lobatchevsky Nizhny Novgorod State University
23 Gagarin ave., 603600 Nizhny Novgorod, Russia
e-mail: balandin @ mm.unn.ru

O.N. Pakhareva
Department of Mathematics and Mechanics
N.I.Lobatchevsky Nizhny Novgorod State University
23 Gagarin ave., 603600 Nizhny Novgorod, Russia
e-mail: pakhareva@rambler.ru

The Lax representation and Backlund transformations for the systems similar to WZNW (Wess-Zumino-Novicov-Witten) systems and non-abelian affine Toda models are obtained in present paper. One of these systems is a new integrable extension of well known sine-Gordon equation.

1 Introduction

Chiral-type systems (see e.g. [1]) are the systems of partial differential equations of the form

$$U^{\alpha}_{xy} + G^{\alpha}_{\beta\gamma} U^{\beta}_{x} U^{\gamma}_{y} + Q^{\alpha} = 0. \tag{1}$$

Here the Greek indices α, β, γ range from 1 to n and the subscripts denote partial derivatives with respect to the independent variables x and y. The coefficients $G^{\alpha}_{\beta\gamma}, Q^{\alpha}$ are assumed to be smooth functions of the variables $U^1, U^2, ..., U^n$. The summation rule over the repeated indices is also assumed.

If system (1) is the system of Euler-Lagrange equations then the Lagrangian can be given by

$$L = g_{\alpha\beta}(U^{\delta}) U^{\alpha}_{x} U^{\beta}_{y} + a_{\alpha\beta}(U^{\delta}) U^{\alpha}_{x} U^{\beta}_{y} + Q(U^{\delta}), \tag{2}$$

where $g_{\alpha\beta}$ is a non-degenerate symmetric matrix, $a_{\alpha\beta}$ is a skew-symmetric matrix and Q is a smooth function of the variables $U^1, U^2, ..., U^n$. Notice that

---

[1] corresponding author

Euler-Lagrange systems for Lagrangian (2) usually are referred as general nonlinear sigma-models (see e.g. [2] ,[3]). These systems contain the special class of systems referred as WZNW(Wess-Zumino-Novicov-Witten)([4] ,[5] , [6], [7]).

Recall that the WZNW systems can be defined in the following way. Let H be a semi-simple Lie group, (U^α) a local coordinate system on H and $\Phi^\alpha = T^\alpha_\beta dU^\beta$ a basis of left-invariant forms. Then

$$G^\alpha_{\beta\gamma} = \tilde{T}^\alpha_\delta T^\delta_{\beta,\gamma} \tag{3}$$

are the coefficients of the first flat affine connection of the group H. Here and further $\tilde{T}^\alpha_\beta$ denotes the inverse matrix for T^α_β and comma denotes the partial derivative

$$T^\delta_{\beta,\gamma} = \frac{\partial T^\delta_\beta}{\partial U^\gamma}.$$

WZNW system associated with the semi-simple Lie group H is the system of the form (1) with $Q^\alpha = 0$ and coefficients $G^\alpha_{\beta\gamma}$ defined by (3).

To obtain the Lagrangian L_1 for WZNW system define a 2-form $\sigma = a_{\alpha\beta} dU^\alpha \wedge dU^\beta$ which satisfies the condition

$$d\sigma = \frac{2}{3} h^0_{\alpha\delta} C^\delta_{\beta\gamma} \Phi^\alpha \wedge \Phi^\gamma \wedge \Phi^\beta.$$

Here $C^\alpha_{\beta\gamma}$ are the structure constants of the group H determined by the basis Φ^α and $h^0_{\alpha\beta}$ is the Killing form of the Lie algebra $\mathfrak{h}$ of the group H with respect to the dual basis. Such 2-form σ exist, at least locally, since 3-form $\Psi = \frac{2}{3} h^0_{\sigma\delta} C^\sigma_{\varphi\psi} T^\delta_\alpha T^\varphi_\beta T^\psi_\gamma dU^\alpha \wedge dU^\gamma \wedge dU^\beta$ is closed due to the Jacobi identity. Then the Lagrangian L_1 is of the form

$$L_1 = h^0_{\gamma\delta} T^\gamma_\alpha T^\delta_\beta U^\alpha_x U^\beta_y + a_{\alpha\beta} U^\alpha_x U^\beta_y. \tag{4}$$

It is well known that WZNW systems admit the Lax representation.

In Section 2 the Euler-Lagrange systems for the Lagrangian

$$L_2 = h^0_{\gamma\delta} T^\gamma_\alpha T^\delta_\beta U^\gamma_x U^\delta_y + k a_{\alpha\beta} U^\alpha_x U^\beta_y$$

are considered. Here k is an arbitrary constant, $h^0_{\alpha\beta}, T^\alpha_\beta, a_{\alpha\beta}$ are of the same sense as above. We will call such systems the WZNW-like systems. The Lax representation and Backlund transformation for these WZNW-like systems are constructed in Section 2.

In Section 3 we give a slight addition to Lesnov-Saveliev construction [8] of Lax representation of nonlinear systems. Lesnov-Saveliev method affords to associate some integrable system with symmetric space G/H. The systems obtained by this way are usually referred as non-abelian affine Toda models ([3] , [9]). Our observation affords to obtain additional integrable systems in case H is a direct product of simple Lie groups. Further we mention some new integrable systems associated with the symmetric spaces $SO(p+3)/(SO(p)\times SO(3)), SO(p+2)/(SO(p)\times SO(2))$.

2 WZNW-like systems

A construction of the Lax representation for some chiral field systems

$$U^{\alpha}_{xy} + G^{\alpha}_{\beta\gamma} U^{\beta}_{x} U^{\gamma}_{y} = 0 \tag{5}$$

is proposed here.

Let Θ^a be a basis of left-invariant forms of a Lie group G. It is well known that the forms Θ^a satisfy the Cartan structure equations

$$d\Theta^a = C^a_{bc}\Theta^c \wedge \Theta^b, \tag{6}$$

where C^a_{bc} are the structure constants.

We say that system (5) admits a Lax representation with the Lie group G if the system

$$\Theta^a = A^a dx + B^a dy \tag{7}$$

is completely integrable (in the sense of Frobenius) on solutions of the system (5), i.e. if the substitution of expressions (7) into (6) results in the system equivalent to (5).

We will consider the groups for which Cartan's structure equations can be written in the form

$$d\theta^{\alpha} = D^{\alpha}_{\beta i}\varphi^i \wedge \theta^{\beta}, \tag{8}$$

$$d\varphi^i = C^i_{jk}\varphi^k \wedge \varphi^j + R^i_{\beta\gamma}\theta^{\gamma} \wedge \theta^{\beta}, \tag{9}$$

where $D^{\alpha}_{\beta i}, C^i_{jk}, R^i_{\beta\gamma} = const$, the Greece indices α, β, γ range from 1 to n (n=dimG/H), indices i, j, k range from 1 to r.

It is well known [10] that Eqs.(8) and (9) could also be considered as the structure equations of some locally symmetric space G/H, where C^i_{jk} are the structure constants of the isotropy group H.

We look for the Lax representation of system (5) in the form

$$\theta^\alpha = U^\alpha_\beta(\lambda U^\beta_x dx + \frac{1}{\lambda} U^\beta_y dy), \tag{10}$$

$$\varphi^i = (-\Gamma^i_\alpha + H^i_\alpha) U^\alpha_\beta U^\beta_x dx + (-\Gamma^i_\alpha - H^i_\alpha) U^\alpha_\beta U^\beta_y dy, \tag{11}$$

where λ is a parameter, $U^\alpha_\beta, \Gamma^i_\alpha, H^i_\alpha$ are the smooth functions of variables $U^1, U^2, ..., U^n$. Substitute the expressions (10),(11) into Eqs.(8). Taking into account (5) and collecting similar terms, we arrive at equations

$$U^\alpha_{\beta,\gamma} - U^\alpha_\delta G^\delta_{\beta\gamma} + D^\alpha_{\mu i}(\Gamma^i_\nu + H^i_\nu) U^\mu_\beta U^\nu_\gamma = 0, \tag{12}$$

$$U^\alpha_{\gamma,\beta} - U^\alpha_\delta G^\delta_{\beta\gamma} + D^\alpha_{\mu i}(\Gamma^i_\nu - H^i_\nu) U^\mu_\gamma U^\nu_\beta = 0. \tag{13}$$

Subtracting (13) from (12) and separating out the symmetric and skew-symmetric parts of result, we obtain the equalities

$$U^\alpha_{[\beta,\gamma]} + D^\alpha_{\mu i} \Gamma^i_\nu U^\mu_{[\beta} U^\nu_{\gamma]} = 0, \tag{14}$$

$$D^\alpha_{\mu i} H^i_\nu U^\mu_{(\beta} U^\nu_{\gamma)} = 0. \tag{15}$$

Now from Eqs.(12),(13) in terms of (14),(15) it follows that

$$U^\alpha_\delta G^\delta_{[\beta\gamma]} = D^\alpha_{\mu i} H^i_\nu U^\mu_{[\beta} U^\nu_{\gamma]}, \tag{16}$$

$$U^\alpha_\delta G^\delta_{(\beta\gamma)} = U^\alpha_{(\beta,\gamma)} + D^\alpha_{\mu i} \Gamma^i_\nu U^\mu_{(\beta} U^\nu_{\gamma)}. \tag{17}$$

Analogously, substituting (10) and (11) into Eqs.(9) and taking into account (5),(14),(15),(16),(17), we get

$$\Gamma^i_{[\mu;\nu]} + H^i_\sigma D^\sigma_{[\mu|j|} H^j_{\nu]} - \Gamma^i_\sigma D^\sigma_{[\mu|j|} \Gamma^j_{\nu]} = C^i_{jk} \Gamma^k_{[\mu} \Gamma^j_{\nu]} - C^i_{jk} H^k_{[\mu} H^j_{\nu]} - R^i_{\mu\nu}, \tag{18}$$

$$H^i_{(\mu;\nu)} = H^i_\sigma D^\sigma_{(\mu|j|} \Gamma^j_{\nu)} + 2 C^i_{jk} H^k_{(\mu} \Gamma^j_{\nu)}. \tag{19}$$

Here we assume $d\Gamma^i_\nu = \Gamma^i_{\nu;\sigma} U^\sigma_\beta dU^\beta$, $dH^i_\nu = H^i_{\nu;\sigma} U^\sigma_\beta dU^\beta$. So the following theorem is proved.

Theorem 1 *Let G be a Lie group with the structure equations* (8),(9) *and* $G^{\alpha}_{\beta\gamma}, U^{\alpha}_{\beta}, \Gamma^{i}_{\alpha}, H^{i}_{\alpha}$ *are smooth functions satisfying the conditions* (14)-(19). *Then equalities* (10),(11) *define the Lax representation of the system* (5).

Example 1.

The proposed construction can be used to obtain well known Lax representation of chiral fields equations valued in symmetric space G/H (see e.g. [11]).

Let $\theta^{\alpha} = U^{\alpha}_{\beta} dU^{\beta}$ are the basic forms and $\varphi^{i} = -\Gamma^{i}_{\alpha} U^{\alpha}_{\beta} dU^{\beta}$ are the connection forms of the symmetric space G/H. Then from (8),(9) it follows that

$$U^{\alpha}_{[\mu,\nu]} = -D^{\alpha}_{\varphi i} \Gamma^{i}_{\psi} U^{\varphi}_{[\mu} U^{\psi}_{\nu]},$$

$$\Gamma^{i}_{[\mu,\nu]} - \Gamma^{i}_{\sigma} D^{\sigma}_{[\mu|i|} \Gamma^{i}_{\nu]} = C^{i}_{jk} \Gamma^{k}_{\mu} \Gamma^{j}_{\nu} - R^{i}_{\mu\nu}.$$

Assume $H^{i}_{\alpha} = 0$ in (11). Then the conditions (14),(15),(18),(19) are fulfilled and equations(8)-(11) define the Lax representation of the system (5), where

$$G^{\alpha}_{[\beta\gamma]} = 0,$$

$$G^{\alpha}_{(\beta\gamma)} = \tilde{U}^{\alpha}_{\sigma} [U^{\sigma}_{(\beta,\gamma)} + D^{\alpha}_{\varphi i} \Gamma^{i}_{\psi} U^{\varphi}_{(\beta} U^{\psi}_{\gamma)}],$$

i.e. $G^{\alpha}_{\beta\gamma}$ are the coefficients of the canonical affine connection of the symmetric space G/H with respect to the holonomic basis.

Example 2.

Let H be a Lie group. Consider the symmetric space $(H \times H)/H$ with the canonical affine connection. Then the structure equations of the symmetric space $(H \times H)/H$ are of the form (8),(9), where C^{i}_{jk} are the structure constants of the group H, $D^{\alpha}_{\beta i} = 2C^{\alpha}_{\beta i}$, $R^{i}_{\alpha\beta} = -C^{i}_{\alpha\beta}$ and all indices range from 1 to $n = dim H$. Assuming $\Gamma^{i}_{\alpha} = 0, H^{i}_{\alpha} = \delta^{i}_{\alpha}$ in (10),(11), we obtain the Lax representation of the system (5) with coefficients $G^{\alpha}_{\beta\gamma}$ of the form

$$G^{\alpha}_{[\beta\gamma]} = 2\tilde{U}^{\alpha}_{\delta} C^{\delta}_{\varphi\psi} U^{\varphi}_{\beta} U^{\psi}_{\gamma},$$

$$G^{\alpha}_{(\beta\gamma)} = 2\tilde{U}^{\alpha}_{\delta} U^{\delta}_{\beta,\gamma}.$$

Here U^{α}_{β} are the smooth functions with the following properties

$$U^{\alpha}_{[\beta,\gamma]} = 0,$$

$$det||U^{\alpha}_{\beta}|| \neq 0.$$

Theorem 2 *Let H be a semi-simple Lie group and*
1) (U^α) is a local coordinate system on the group H;
2) $C^\alpha_{\beta\gamma}$ are the structure constants of the Lie group H with respect to the basis of left-invariant forms $\Phi^\alpha = T^\alpha_\beta dU^\beta$;
3) $h^0_{\alpha\beta}$ is the Killing form of the Lie algebra $\mathfrak{h}$ of the group H given in the dual basis;
4) $h_{\alpha\beta}$ is the Killing metric of the group H, i.e.

$$h_{\alpha\beta} = h^0_{\varphi\psi} T^\varphi_\alpha T^\psi_\beta; \tag{20}$$

5) $\sigma = a_{\alpha\beta} dU^\alpha \wedge dU^\beta$ is the 2-form which satisfies the condition

$$d\sigma = \frac{2}{3} h^0_{\delta\sigma} C^\sigma_{\beta\gamma} \Phi^\delta \wedge \Phi^\gamma \wedge \Phi^\beta.$$

Then the system of Euler-Lagrange equations for Lagrangian

$$L = h_{\alpha\beta} U^\alpha_x U^\beta_y + k a_{\alpha\beta} U^\alpha_x U^\beta_y \tag{21}$$

admits the Lax representation. Here $k \neq \pm 1$ is an arbitrary constant.

Proof. Consider the symmetric space $(H \times H)/H$ with the structure equations (8),(9), where $D^\alpha_{\beta i} = 2C^\alpha_{\beta i}$, $R^i_{\alpha\beta} = -C^i_{\alpha\beta}$ and all indices range from 1 to n. Assume

$$U^\alpha_\beta = \frac{1}{2}\sqrt{k^2-1}\, T^\alpha_\beta,$$

$$\Gamma^i_\alpha = -\frac{1}{\sqrt{k^2-1}}\delta^i_\alpha, H^i_\alpha = \frac{k}{\sqrt{k^2-1}}\delta^i_\alpha,$$

where $k \neq \pm 1$ is an arbitrary constant. Then the expressions (10),(11) define the Lax representation for the system

$$U^\alpha_{xy} + \tilde{T}^\alpha_\sigma [T^\sigma_{(\beta,\gamma)} + k C^\sigma_{\varphi\psi} T^\varphi_\beta T^\psi_\gamma] U^\beta_x U^\gamma_y = 0. \tag{22}$$

Now prove that the system (22) is the system of Euler's equations for Lagrangian (21). Indeed, the Euler equations for the Lagrangian (21) results in the system (5), where

$$G^\alpha_{\beta\gamma} = \frac{1}{2}\tilde{h}^{\alpha\delta}(h_{\delta\gamma,\beta} + h_{\delta\beta,\gamma} - h_{\beta\gamma,\delta}) + \frac{3}{2}\tilde{h}^{\alpha\delta} a_{[\delta\gamma,\beta]}. \tag{23}$$

One can easily verify that in our case we have the equalities

$$G^{\alpha}_{(\beta\gamma)} = \frac{1}{2}\tilde{h}^{\alpha\delta}(h_{\delta\gamma,\beta} + h_{\delta\beta,\gamma} - h_{\beta\gamma,\delta}) = \tilde{T}^{\alpha}_{\delta} T^{\delta}_{(\beta,\gamma)}. \tag{24}$$

So the coefficients $G^{\alpha}_{\beta\gamma}$ of the Euler equations are of the form (22) if the torsion

$$A^{\alpha}_{\beta\gamma} = G^{\alpha}_{[\beta\gamma]} = k\tilde{T}^{\alpha}_{\sigma} C^{\sigma}_{\varphi\psi} T^{\varphi}_{\beta} T^{\psi}_{\gamma} \tag{25}$$

of the connection satisfies the following condition $A_{\alpha\beta\gamma} = h_{\delta\alpha} A^{\delta}_{\beta\gamma} = \frac{3}{2} a_{[\alpha\gamma,\beta]}$. Taking into account (20), we obtain $A_{\alpha\beta\gamma} = kh^{0}_{\sigma\delta} C^{\sigma}_{\varphi\psi} T^{\delta}_{\alpha} T^{\varphi}_{\beta} T^{\psi}_{\gamma}$.

Notice that 3-form $\Psi = \frac{2}{3} h^{0}_{\sigma\delta} C^{\sigma}_{\varphi\psi} T^{\delta}_{\alpha} T^{\varphi}_{\beta} T^{\psi}_{\gamma} dU^{\alpha} \wedge dU^{\gamma} \wedge dU^{\beta}$ is closed due to the Jacobi identity. Therefore there exist, at least locally, a 2-form σ such that the equality $d\sigma = \Psi$ is fulfilled. So assuming $a_{\alpha\beta} dU^{\alpha} \wedge dU^{\beta} = \sigma$, we obtain that the Euler equations for Lagrangian (21) coincide with the system (22). $\Box$

Remark. One can easily verify that the coefficients $G^{\alpha}_{\beta\gamma}$ defined by equalities (24),(25) are the coefficients of the affine connection of homogeneous reductive space which can be characterized by the following conditions:

$$\nabla R^{\alpha}_{\beta\gamma\delta} = 0, \ \nabla A^{\alpha}_{\beta\gamma} = 0.$$

Moreover, one can see that the Ricci identities

$$R^{\alpha}_{[\beta\gamma\delta]} = 0$$

are true for the connection with coefficients defined by (24),(25).

Notice that the Lax representation without parameter for the systems with such connection was mentioned in [12].

The Backlund transformation for the systems (22) can be obtained from the following theorem

Theorem 3 *Let $\overline{C}^{\alpha}_{\beta\gamma}, \widehat{C}^{\alpha}_{\beta\gamma}$ be the structure constants of n-dimensional Lie algebras and functions $U^{\alpha}_{\beta} = U^{\alpha}_{\beta}(U^{\gamma}), V^{\alpha}_{\beta} = V^{\alpha}_{\beta}(V^{\gamma})$ satisfy the equations:*

$$U^{\alpha}_{[\beta,\gamma]} = \overline{C}^{\alpha}_{\varphi\psi} U^{\varphi}_{\beta} U^{\psi}_{\gamma}, \ \ det||U^{\alpha}_{\beta}|| \neq 0, \tag{26}$$

$$V^{\alpha}_{[\beta,\gamma]} = \widehat{C}^{\alpha}_{\varphi\psi} V^{\varphi}_{\beta} V^{\psi}_{\gamma}, \ \ det||V^{\alpha}_{\beta}|| \neq 0, \tag{27}$$

Then the equalities

$$U^\alpha_\beta U^\beta_x = V^\alpha_\beta V^\beta_x, \tag{28}$$

$$U^\alpha_\beta U^\beta_y = -V^\alpha_\beta V^\beta_y \tag{29}$$

define the Backlund transformations between the systems

$$U^\alpha_{xy} + \tilde{U}^\alpha_\delta [U^\delta_{(\beta,\gamma)} + \widehat{C}^\delta_{\varphi\psi} U^\varphi_\beta U^\psi_\gamma] U^\beta_x U^\gamma_y = 0, \tag{30}$$

$$V^\alpha_{xy} + \tilde{V}^\alpha_\delta [V^\delta_{(\beta,\gamma)} + \overline{C}^\delta_{\varphi\psi} V^\varphi_\beta V^\psi_\gamma] V^\beta_x V^\gamma_y = 0 \tag{31}$$

Proof. Indeed, take the derivatives of equalities (28),(29) with respect to variables x and y. Then adding and subtracting the obtained equalities and taking into account (26),(27),(28),(29), we arrive at the systems (30),(31). □

Corollary 1. *The equalities* (28),(29) *define the Backlund transformations between the systems*

$$U^\alpha_{xy} + \tilde{U}^\alpha_\delta [U^\delta_{(\beta,\gamma)} + k C^\delta_{\varphi\psi} U^\varphi_\beta U^\psi_\gamma] U^\beta_x U^\gamma_y = 0, \tag{32}$$

$$V^\alpha_{xy} + \tilde{W}^\alpha_\delta [W^\delta_{(\beta,\gamma)} + \frac{1}{k} C^\delta_{\varphi\psi} W^\varphi_\beta W^\psi_\gamma] V^\beta_x V^\gamma_y = 0, \tag{33}$$

were $k = const$ *and functions* $U^\alpha_\beta = U^\alpha_\beta(U^\gamma), W^\alpha_\beta = W^\alpha_\beta(V^\gamma)$ *satisfy the equations*

$$U^\alpha_{[\beta,\gamma]} = C^\alpha_{\varphi\psi} U^\varphi_\beta U^\psi_\gamma, \;\; det||U^\alpha_\beta|| \neq 0, \tag{34}$$

$$W^\alpha_{[\beta,\gamma]} = C^\alpha_{\varphi\psi} W^\varphi_\beta W^\psi_\gamma, \;\; det||W^\alpha_\beta|| \neq 0. \tag{35}$$

Proof. Assume $\overline{C}^\alpha_{\beta\gamma} = C^\alpha_{\beta\gamma}, \widehat{C}^\alpha_{\beta\gamma} = kC^\alpha_{\beta\gamma}, W^\alpha_\beta = kV^\alpha_\beta$. Then the functions U^α_β, W^α_β satisfy the Eqs.(35),(34) and the systems (30),(31) coincide with the systems (32),(33).□

3 Systems associated with the symmetric space $G/(H_1 \times ... \times H_p)$

We use the local coordinate approach to Lesnov-Saveliev construction in order to find the Lax representation for some systems of the form (1).

Let G/H be a symmetric space with the structure equations of the form

$$d\omega^{\alpha'} = D^{\alpha'}_{\beta'\gamma}\theta^{\gamma} \wedge \omega^{\gamma'}, \tag{36}$$

$$d\theta^{\alpha} = C^{\alpha}_{\beta\gamma}\theta^{\gamma} \wedge \theta^{\beta} + R^{\alpha}_{\beta'\gamma'}\omega^{\gamma'} \wedge \omega^{\beta'}. \tag{37}$$

Here the Greek indices α, β, γ range from 1 to n and indices α', β', γ' range from $n+1$ to r. We assume

$$\omega^{\alpha'} = \lambda M^{\alpha'} dx + \frac{1}{\lambda} N^{\alpha'} dy, \tag{38}$$

$$\theta^{\alpha} = T^{\ \alpha}_{1\beta} U^{\beta}_{x} dx + T^{\ \alpha}_{2\beta} U^{\beta}_{y} dy, \tag{39}$$

where $M^{\alpha'}, N^{\alpha'}, T^{\ \alpha}_{1\beta}, T^{\ \alpha}_{2\beta}$ are the smooth functions of variables $U^1, ..., U^n$.One can easily verify that the following theorem is true [13].

Theorem 4 *Let* $C^{\alpha}_{\beta\gamma}$ *be the structure constants of the isotropy group* H *of some locally symmetric space* G/H *with the structure equations* (36),(37). *Assume that there exist matrices* $\|T^{\ \alpha}_{1\beta}\|, \|T^{\ \alpha}_{2\beta}\|$ *and functions* $M^{\alpha'}, N^{\alpha'}$ *with the following properties:*

$$T^{\ \alpha}_{i[\beta,\gamma]} = C^{\alpha}_{\mu\nu} T^{\ \mu}_{i\beta} T^{\ \nu}_{i\gamma} \ (i = 1, 2), \tag{40}$$

$$det\|T^{\ \alpha}_{1\beta} - T^{\ \alpha}_{2\beta}\| \neq 0, \tag{41}$$

$$M^{\alpha'}_{,\delta} = D^{\alpha'}_{\beta'\gamma} T^{\ \gamma}_{2\delta} M^{\beta'}, \tag{42}$$

$$N^{\alpha'}_{,\delta} = D^{\alpha'}_{\beta'\gamma} T^{\ \gamma}_{1\delta} N^{\beta'}. \tag{43}$$

Then equalities (38),(39) *define the Lax representation of the system* (1) *if the functions* $G^{\alpha}_{\beta\gamma}, Q^{\alpha}$ *are of the form*

$$G^{\alpha}_{\beta\gamma} = \tilde{P}^{\alpha}_{\delta}[P^{\delta}_{(\beta,\gamma)} + 2C^{\delta}_{\varphi\psi} S^{\varphi}_{\beta} S^{\psi}_{\gamma} - 2C^{\delta}_{\varphi\psi} P^{\varphi}_{(\gamma} S^{\psi}_{\beta)}],$$

$$Q^\alpha = -\tilde{P}^\alpha_\delta R^\delta_{\beta'\gamma'} N^{\gamma'} M^{\beta'},$$

where $S^\alpha_\beta = \frac{1}{2}(T^{\ \alpha}_{1\beta} + T^{\ \alpha}_{2\beta})$, $P^\alpha_\beta = \frac{1}{2}(T^{\ \alpha}_{1\beta} - T^{\ \alpha}_{2\beta})$ *and* $\tilde{P}$ *denotes the inverse matrix for* P*.*

The question under consideration is how to find the functions $T^{\ \alpha}_{i\beta}, M^{\alpha'}, N^{\alpha'}$ satisfying the condition (40)-(43). One solution can be obtained by the following way. Let $C^\alpha_{\beta\gamma}$ be the structure constants of the group H with respect to a basis of left-invariant forms $\Phi^\alpha = T^\alpha_\beta dU^\beta$. Assume $T^{\ \alpha}_{1\beta} = T^\alpha_\beta,\ T^{\ \alpha}_{2\beta} = 0$ (or $T^{\ \alpha}_{1\beta} = 0,\ T^{\ \alpha}_{2\beta} = T^\alpha_\beta$). Then one can see that (40),(41) and the compatibility conditions for the systems (42),(43) are fulfilled. So the equalities (36)-(39) define the Lax representation of the system (1) with the coefficients Q^α and $G^\alpha_{\beta\gamma}$ mentioned in theorem 4. Notice that this Lax representation coincide with one obtained by Lesnov-Saveliev [8]. Further we give another solution of the equations (40)-(43) in case H is a direct product of simple groups $H_1, ..., H_p$.

Let G/H be a locally symmetric space and $H = H_1 \times H_2$, where H_1, H_2 are the simple Lie groups. Split the local coordinates (U^α) into to groups $(U^{\alpha_1}, U^{\alpha_2})$ according to the decomposition of the group H and rewrite the structure equations of the symmetric space G/H in the form

$$d\omega^{\beta'} = D^{\beta'}_{\gamma'\alpha_1}\theta^{\alpha_1} \wedge \omega^{\gamma'} + D^{\beta'}_{\gamma'\alpha_2}\theta^{\alpha_2} \wedge \omega^{\gamma'}, \tag{44}$$

$$d\theta^{\alpha_1} = C^{\alpha_1}_{\beta_1\gamma_1}\theta^{\gamma_1} \wedge \theta^{\beta_1} + R^{\alpha_1}_{\beta'\gamma'}\omega^{\gamma'} \wedge \omega^{\beta'}, \tag{45}$$

$$d\theta^{\alpha_2} = C^{\alpha_2}_{\beta_2\gamma_2}\theta^{\gamma_2} \wedge \theta^{\beta_2} + R^{\alpha_2}_{\beta'\gamma'}\omega^{\gamma'} \wedge \omega^{\beta'}. \tag{46}$$

We will use the following notations.
1) $\mathfrak{g}, \mathfrak{h}, \mathfrak{h}_1, \mathfrak{h}_2$ are the Lie algebras of the groups Lie G, H, H_1, H_2 respectively.
2) $C^{\alpha_1}_{\beta_1\gamma_1}, C^{\alpha_2}_{\beta_2\gamma_2}$ are the structure constants of the Lie groups H_1, H_2 with respect to the bases of left-invariant forms $\Phi^{\alpha_1} = T^{\alpha_1}_{\beta_1}dU^{\beta_1}, \Phi^{\alpha_2} = T^{\alpha_2}_{\beta_2}dU^{\beta_2}$.
3) $h^0_{\alpha_1\beta_1}, h^0_{\alpha_2\beta_2}$ are the matrices of the Killing forms $h^0_1(\cdot,\cdot), h^0_2(\cdot,\cdot)$ of the Lie algebras $\mathfrak{h}_1, \mathfrak{h}_2$ given in the bases dual to the bases $\Phi^{\alpha_1}, \Phi^{\alpha_2}$ and $g^0(\cdot,\cdot)$ is the Killing form of the algebra $\mathfrak{g}$.
4) S_i are the constants satisfying the equalities

$$h^0_i(\cdot,\cdot) = S_i g^0(\cdot,\cdot)_{|\mathfrak{h}_i} \quad (i = 1, 2). \tag{47}$$

Such constants exist due to simplicity of the subalgebras $\mathfrak{h}_1, \mathfrak{h}_2$.
5) $h_{\alpha_1\beta_1}, h_{\alpha_2\beta_2}$ are the Killing metrics of the groups H_1 and H_2 respectively, i.e.

$$h_{\alpha_i\beta_i} = h^0_{\gamma_i\delta_i} T^{\gamma_i}_{\alpha_i} T^{\delta_i}_{\beta_i}, \quad (i = 1, 2). \tag{48}$$

6) $\sigma_i = a_{\alpha_i\beta_i} dU^{\alpha_i} \wedge dU^{\beta_i}$ $(i = 1, 2)$ are the 2-forms which satisfy the conditions

$$d\sigma_i = \frac{2}{3} h^0_{\alpha_i\delta_i} C^{\delta_i}_{\beta_i\gamma_i} \Phi^{\alpha_i} \wedge \Phi^{\gamma_i} \wedge \Phi^{\beta_i}. \tag{49}$$

Theorem 5 *Let G/H be a locally symmetric space, G the semi-simple Lie group and $H = H_1 \times H_2$, where H_1, H_2 are the simple Lie groups.*

Then the Euler's system for Lagrangian

$$L = S_2[h_{\alpha_1\beta_1}(U^{\gamma_1}) + \varepsilon_1 a_{\alpha_1\beta_1}(U^{\gamma_1})] U^{\alpha_1}_x U^{\beta_1}_y +$$

$$+ S_1[h_{\alpha_2\beta_2}(U^{\gamma_2}) + \varepsilon_2 a_{\alpha_2\beta_2}(U^{\gamma_2})] U^{\alpha_2}_x U^{\beta_2}_y + Q \tag{50}$$

admits the Lax representation if

$$Q = 4 S_1 S_2 g^0_{\gamma'\delta'} M^{\gamma'} N^{\delta'}, \tag{51}$$

$$\varepsilon_i = \pm 1 \; (i = 1, 2),$$

and functions $M^{\gamma'}, N^{\delta'}$ satisfy the following conditions:

$$M^{\gamma'}_{,\beta_1} = \frac{1 - \varepsilon_1}{2} D^{\gamma'}_{\delta'\alpha_1} T^{\alpha_1}_{\beta_1} M^{\delta'}, \quad M^{\gamma'}_{,\beta_2} = \frac{1 - \varepsilon_2}{2} D^{\gamma'}_{\delta'\alpha_2} T^{\alpha_2}_{\beta_2} M^{\delta'}, \tag{52}$$

$$N^{\gamma'}_{,\beta_1} = \frac{1 + \varepsilon_1}{2} D^{\gamma'}_{\delta'\alpha_1} T^{\alpha_1}_{\beta_1} N^{\delta'}, \quad N^{\gamma'}_{,\beta_2} = \frac{1 + \varepsilon_2}{2} D^{\gamma'}_{\delta'\alpha_2} T^{\alpha_2}_{\beta_2} N^{\delta'}. \tag{53}$$

Proof. Assume

$$\theta^{\alpha_1} = \frac{1 + \varepsilon_1}{2} T^{\alpha_1}_{\beta_1} U^{\beta_1}_x dx + \frac{1 - \varepsilon_1}{2} T^{\alpha_1}_{\beta_1} U^{\beta_1}_y dy, \tag{54}$$

$$\theta^{\alpha_2} = \frac{1 + \varepsilon_2}{2} T^{\alpha_2}_{\beta_2} U^{\beta_2}_x dx + \frac{1 - \varepsilon_2}{2} T^{\alpha_2}_{\beta_2} U^{\beta_2}_y dy, \tag{55}$$

$$\omega^{\gamma'} = \lambda M^{\gamma'} dx + \frac{1}{\lambda} N^{\gamma'} dy. \tag{56}$$

Substituting the expressions (54)-(56) into the structure equations (44)-(46), we arrive at the system of the form (1) with coefficients

$$G^{\alpha_1}_{\beta_1\gamma_1} = \tilde{T}^{\alpha_1}_{\delta_1}(T^{\delta_1}_{(\beta_1,\gamma_1)} + \varepsilon_1 T^{\delta_1}_{[\beta_1,\gamma_1]}), \tag{57}$$

$$G^{\alpha_2}_{\beta_2\gamma_2} = \tilde{T}^{\alpha_2}_{\delta_2}(T^{\delta_2}_{(\beta_2,\gamma_2)} + \varepsilon_2 T^{\delta_2}_{[\beta_2,\gamma_2]}), \tag{58}$$

$$Q^{\alpha_1} = 2\varepsilon_1 \tilde{T}^{\alpha_1}_{\delta_1} R^{\delta_1}_{\beta'\gamma'} M^{\gamma'} N^{\beta'}, \tag{59}$$

$$Q^{\alpha_2} = 2\varepsilon_2 \tilde{T}^{\alpha_2}_{\delta_2} R^{\delta_2}_{\beta'\gamma'} M^{\gamma'} N^{\beta'}. \tag{60}$$

Prove that this system is the Euler-Lagrange system for the Lagrangian (50). Indeed, the Euler equations for the Lagrangian (50) results in the system

$$U^{\alpha_1}_{xy} + G^{\alpha_1}_{\beta_1\gamma_1} U^{\beta_1}_x U^{\gamma_1}_y - \frac{1}{2S_2}\tilde{h}^{\alpha_1\beta_1} Q_{,\beta_1} = 0,$$

$$U^{\alpha_2}_{xy} + G^{\alpha_2}_{\beta_2\gamma_2} U^{\beta_2}_x U^{\gamma_2}_y - \frac{1}{2S_1}\tilde{h}^{\alpha_2\beta_2} Q_{,\beta_2} = 0,$$

where the coefficients $G^{\alpha_1}_{\beta_1\gamma_1}, G^{\alpha_2}_{\beta_2\gamma_2}$ are of the form (57),(58). The proof of this fact is analogous to the proof of the theorem 2.

Now prove that the functions $-\frac{1}{2S_2}\tilde{h}^{\alpha_1\beta_1}Q_{,\beta_1}$ coincide with the functions (59). Let $\mathfrak{g} = \mathfrak{h} \oplus \mathfrak{m}$ be a canonical decomposition, i.e.

$$[\mathfrak{h},\mathfrak{h}] \subset \mathfrak{h}, [\mathfrak{h},\mathfrak{m}] \subset \mathfrak{m}, [\mathfrak{m},\mathfrak{m}] \subset \mathfrak{h}.$$

Then due to the properties of the Killing form and (47) it is true that

$$h^0_1(h, [m_1, m_2]) = S_1 g^0(m_1, [m_2, h]) \tag{61}$$

for any arbitrary elements $h \in \mathfrak{h}_1,\ \ m_1, m_2 \in \mathfrak{m}$.

The equalities (61) are equivalent to the following conditions

$$h^0_{\alpha_1\varphi_1} R^{\varphi_1}_{\beta'\gamma'} = S g^0_{\beta'\delta'} D^{\delta'}_{\gamma'\alpha_1}. \tag{62}$$

Now differentiating expression (51) and taking into account (52),(53) and (62), we obtain

$$Q_{,\beta_1} = 4S_1 S_2 g^0_{\gamma'\delta'}(M^{\gamma'}_{,\beta_1} N^{\delta'} + M^{\gamma'} N^{\delta'}_{,\beta_1}) =$$

$$= 4S_1S_2g^0_{\gamma'\delta'}(\frac{1-\varepsilon_1}{2}D^{\gamma'}_{\psi'\alpha_1}T^{\alpha_1}_{\beta_1}M^{\psi'}N^{\delta'} + \frac{1+\varepsilon_1}{2}D^{\delta'}_{\psi'\alpha_1}T^{\alpha_1}_{\beta_1}M^{\gamma'}N^{\psi'}) =$$

$$= -4\varepsilon_1 S_2 h^0_{\alpha_1\varphi_1} R^{\varphi_1}_{\delta'\gamma'} T^{\alpha_1}_{\beta_1} M^{\gamma'} N^{\delta'}$$

and the functions $-\frac{1}{2S_2}\tilde{h}^{\alpha_1\beta_1}Q_{,\beta_1}$ coincide with the functions (59). Analogously one can verify that the functions $-\frac{1}{2S_1}\tilde{h}^{\alpha_2\beta_2}Q_{,\beta_2}$ coincide with the functions (60).$\Box$

Remark. In case $\varepsilon_1 = \varepsilon_2$ we come to the system which can be obtained by means of Lesnov-Saveliev construction.

Analogously can be proved the following theorem.

Theorem 6 *Let G/H be a locally symmetric space, G the semi-simple Lie group and $H = H_1 \times ... \times H_p$, where $H_1, ..., H_p$ are the simple Lie groups.*

Then the Euler's system for Lagrangian

$$L = \sum_{i=1}^{p} \frac{S}{S_i}[h_{\alpha_i\beta_i}(U^{\gamma_i}) + \varepsilon_i a_{\alpha_i\beta_i}(U^{\gamma_i})]U^{\alpha_i}_x U^{\beta_i}_y + 4Sg^0_{\beta'\gamma'}M^{\beta'}N^{\gamma'}$$

admits the Lax representation. Here $\varepsilon_i = \pm 1$; $h_{\alpha_i\beta_i}$ are the Killing metrics of the groups H_i $(i = \overline{1,p})$; $S = S_1...S_p$; constants S_i are defined by equalities

$$h^0_i(\cdot,\cdot) = S_i g^0(\cdot,\cdot)_{|\mathfrak{h}_i};$$

$a_{\alpha_i\beta_i}$ are defined as in previous theorems by 2-forms σ_i; and $M^{\beta'}, N^{\gamma'}$ are defined analogously to previous theorem.

Example 3.(The system associated with $SO(6)/(SO(3) \times SO(3))$.)

Choose the coordinates U^1, U^2, U^3 of the group $SO(3)$ to be Euler's angles. Then the left-invariant forms can be written in the following form

$$\Phi^1 = -\cos U^2 dU^3 - \sin U^2 \sin U^3 dU^1,$$

$$\Phi^2 = \sin U^3 \cos U^2 dU^1 - \sin U^2 dU^3,$$

$$\Phi^3 = -\cos U^3 dU^1 - dU^2$$

and

$$d\Phi^1 = \Phi^3 \wedge \Phi^2,\ \Phi^2 = \Phi^1 \wedge \Phi^3,\ \Phi^3 = \Phi^2 \wedge \Phi^1$$

are the structure equations of the group $SO(3)$. On the second group $H_2 = SO(3)$ we choose the local coordinates U^4, U^5, U^6 and left-invariant forms Φ^4, Φ^5, Φ^6 analogously.

Rewrite the structure equations of the symmetric space $SO(6)/(SO(3) \times SO(3))$ in the form

$$d\Omega = \Omega \wedge \Omega,\ \Omega = \left\| \begin{array}{cc} \theta^a_b & \omega^a_{b'} \\ \omega^{a'}_b & \theta^{a'}_{b'} \end{array} \right\|, \tag{63}$$

where the forms $\theta^a_b = -\theta^b_a, \theta^{a'}_{b'} = -\theta^{b'}_{a'}, \omega^{a'}_b = -\omega^b_{a'}$, the indices $a, b, c...$ range from 1 to 3 and $a', b', ...$ range from 4 to 6.

Assume $\varepsilon_1 = -1, \varepsilon_2 = 1$, i.e.

$$\theta^1_2 = (-\cos U^3 U^1_y - U^2_y)dy,\ \theta^1_3 = (-\sin U^3 \cos U^2 U^1_y + \sin U^2 U^3_y)dy,$$

$$\theta^2_3 = (-\cos U^2 U^3_y - \sin U^2 \sin U^3 U^1_y)dy,\ \theta^4_5 = (-\cos U^6 U^4_x - U^5_x)dx, \tag{64}$$

$$\theta^4_6 = (-\sin U^6 \cos U^5 U^4_x + \sin U^5 U^6_x)dx,\ \theta^5_6 = (-\cos U^5 U^6_x - \sin U^5 \sin U^6 U^4_x)dx,$$

$$\omega^a_{a'} = \lambda M^a_{a'} dx + \frac{1}{\lambda} N^a_{a'} dy. \tag{65}$$

Substituting the expressions (64),(65) into Eqs.(63) and equating the coefficients with λ and $\frac{1}{\lambda}$, we arrive at the system of partial differential equations for the functions $M^a_{a'}, N^a_{a'}$. Integrating this system, we find the simplest solution: $M^1_6 = \sin U^2 \sin U^3$, $M^2_6 = -\cos U^2 \sin U^3$, $M^3_6 = \cos U^3$, $N^3_4 = k \sin U^2 \sin U^3$, $N^3_5 = -k \cos U^5 \sin U^6$, $N^3_6 = k \cos U^6$, where $k = const$, and the other functions $M^a_{a'}, N^a_{a'}$ vanish. Substituting these functions $M^a_{a'}, N^a_{a'}$, into Eqs.(65), we obtain the Lax representation of the Euler-Lagrange system for Lagrangian

$$L = \sum_{\beta_1=1}^{3} U^{\beta_1}_x U^{\beta_1}_y + 2\cos U^3 U^1_y U^2_x + \sum_{\beta_2=4}^{6} U^{\beta_2}_x U^{\beta_2}_y$$

$$+\cos U^6 (U^4_y U^5_x + U^4_x U^5_y) - \cos U^6 (U^4_y U^5_x - U^4_x U^5_y) + 2k \cos U^3 \cos U^6. \tag{66}$$

Example 4.(An integrable extension of the sine-Gordon equation)

Notice that the system of Euler-Lagrange equations for the Lagrangian (66) admits the reduction which is similar to one shown in [14]. Indeed, inserting the expressions

$$U^1_x = -\cos U^3 U^2_x,\ U^1_y = -\frac{1}{\cos U^3} U^2_y,$$

$$U^4_x = -\frac{1}{\cos U^6} U^5_x,\ U^4_y = -\cos U^6 U^5_y$$

in this system, we arrive at the system

$$U^2_{xy}+U^3_y U^2_x ctg U^3+\frac{1}{\sin U^3 \cos U^3} U^3_x U^2_y = 0,\ U^3_{xy}-U^2_y U^2_x tg U^3+k\sin U^3 \cos U^6 = 0,$$

$$U^5_{xy}+U^5_y U^6_x ctg U^6+\frac{1}{\sin U^6 \cos U^6} U^5_x U^6_y = 0,\ U^6_{xy}-U^5_y U^5_x tg U^6+n\cos U^3 \sin U^6 = 0.$$

The latter system is not the system of Euler-Lagrange equations. However, it admits the Backlund transformation defined by the equalities

$$U^2_y = \frac{\cos U^3}{1+\cos U^3} V^1_y,\quad U^2_x = \frac{1}{1+\cos U^3} V^1_x,\quad U^3 = V^2,$$

$$U^5_y = \frac{1}{1+\cos U^6} V^3_y,\quad U^5_x = \frac{\cos U^6}{1+\cos U^6} V^3_x,\quad U^6 = V^4,$$

to the Euler-Lagrange one

$$\begin{aligned} V^1_{xy} + \frac{1}{\sin V^2} V^1_y V^2_x + \frac{1}{\sin V^2} V^2_y V^1_x &= 0, \\ V^2_{xy} - \frac{\sin V^2}{(1+\cos V^2)^2} V^1_y V^1_x + k\sin V^2 \cos V^4 &= 0, \\ V^3_{xy} + \frac{1}{\sin V^4} V^3_y V^4_x + \frac{1}{\sin V^4} V^4_y V^3_x &= 0, \\ V^4_{xy} - \frac{\sin V^4}{(1+\cos V^4)^2} V^3_y V^3_x + k\cos V^2 \sin V^4 &= 0. \end{aligned} \tag{67}$$

The Lagrangian for the system (67) is of the form

$$L = V^1_x V^1_y tg^2 \frac{V^2}{2} + V^2_x V^2_y + V^3_x V^3_y tg^2 \frac{V^4}{2} + V^4_x V^4_y + 2k\cos V^2 \cos V^4. \tag{68}$$

Remark 1. The system (67) could be considered as a "doubled" complex sine-Gordon system. Indeed, inserting $V^1 = V^3, V^2 = V^4$ in (67), we arrive at the complex sine-Gordon system. Further, assuming $V^1 = V^3 = V,\ V^2 = V^4 = 0$ we obtain well known sine-Gordon equation.

Remark 2. The metric associated with the Lagrangian (68) is the product of two black hole metrics [15].

Remark 3. The system (67) admits the Lax representation which can be written in the form (63), where

$$\theta_2^1 = \frac{1}{2\cos\frac{V^2}{2}}(V_x^1 dx + V_y^1 dy),\ \theta_3^1 = -\frac{\sin\frac{V^2}{2}}{2\cos^2\frac{V^2}{2}}(V_x^1 dx - V_y^1 dy),$$

$$\theta_3^2 = \frac{1}{2}(V_x^2 dx - V_y^2 dy),\ \theta_5^4 = \frac{1}{2\cos\frac{V^4}{2}}(V_x^3 dx + V_y^3 dy),$$

$$\theta_6^4 = -\frac{\sin\frac{V^4}{2}}{2\cos^2\frac{V^4}{2}}(V_x^3 dx - V_y^3 dy),\ \theta_6^5 = \frac{1}{2}(V_x^4 dx - V_y^4 dy),$$

$$\omega_{a'}^a = \lambda M_{a'}^a dx + \frac{1}{\lambda} N_{a'}^a dy,$$

$$M_4^a = M_{a'}^1 = 0,\ N_4^a = N_{a'}^1 = 0,$$

$$M_5^2 = \sin\frac{V^2}{2}\sin\frac{V^4}{2},\ M_6^2 = -\sin\frac{V^2}{2}\cos\frac{V^4}{2},$$

$$M_5^3 = -\cos\frac{V^2}{2}\sin\frac{V^4}{2},\ \ M_6^3 = \cos\frac{V^2}{2}\cos\frac{V^4}{2},$$

$$N_5^2 = k\sin\frac{V^2}{2}\sin\frac{V^4}{2},\ \ N_6^2 = k\sin\frac{V^2}{2}\cos\frac{V^4}{2},$$

$$N_5^3 = k\cos\frac{V^2}{2}\sin\frac{V^4}{2},\ \ N_6^3 = n\cos\frac{V^2}{2}\cos\frac{V^4}{2}.$$

Notice, that we construct the Lax representation of the system (67) with the help of the structure equations of the symmetric space $SO(6)/(SO(3) \times SO(3))$. Further we give examples of new integrable systems similar to Example 4. These systems differ from ones mentioned in theorem 5 but the Lax representations for them are associated with the symmetric spaces of the form $G/(H_1 \times H_2)$.

Theorem 7 *To every symmetric space of the form $SO(p+3)/(SO(p)\times SO(3))$ $(p\geq 3)$ there corresponds the Lax representation of Euler's equations for Lagrangian*

$$L = g_{\alpha_1\beta_1}U_x^{\alpha_1}U_y^{\beta_1} + a_{\alpha_1\beta_1}U_x^{\alpha_1}U_y^{\beta_1} - 2(p-2)[V_x^1V_y^1 tg^2\frac{V^2}{2} + V_x^2V_y^2]$$

$$+4(p-2)M_{b'}^{b}N_{b}^{b'}. \tag{69}$$

Here $g_{\alpha_1\beta_1}, a_{\alpha_1\beta_1}$ are of the same sense as in previous theorems, Latin indices a,b range from 1 to p; indices a',b' range from $p+1$ to $p+3$ and functions $M_{b'}^{b}, N_{b'}^{b}$ will be defined in the proof.

Proof. We use the embedding of the group $SO(p+3)$ into $GL(p+3)$ and rewrite the structure equations of the symmetric space $SO(p+3)/(SO(p)\times SO(3))$ in form (63), where indices $a,b=\overline{1,p},\ a',b'=\overline{p+1,p+3}$. Let the coefficients $H_{b\delta_1}^{a} = -H_{a\delta_1}^{b}$ be defined by the embedding of the Lie algebra $\mathfrak{so}(p)$ into the $\mathfrak{gl}(p)$. Then these coefficients satisfy the identities:

$$H_{b\delta_1}^{a}C_{\beta_1\gamma_1}^{\delta_1} = H_{d[\gamma_1}^{a}H_{|b|\beta_1]}^{d},\ \ (p-2)H_{b\beta_1}^{a}H_{a\gamma_1}^{b} = h_{\beta_1\gamma_1}, \tag{70}$$

where $C_{\beta_1\gamma_1}^{\delta_1}$ and $h_{\beta_1\gamma_1}$ are the structure constants and the Killing metric of $\mathfrak{so}(p)$ whit respect to the basis $\Phi^{\alpha_1} = T_{\beta_1}^{\alpha_1}dU^{\beta_1}$.

Assume

$$\theta_b^a = H_{b\beta_1}^{a}T_{\gamma_1}^{\beta_1}U_x^{\gamma_1}dx,\ \ \theta_{p+2}^{p+1} = \frac{1}{2\cos\frac{V^2}{2}}(V_x^1dx + V_y^1dy), \tag{71}$$

$$\theta_{p+3}^{p+1} = -\frac{\sin\frac{V^2}{2}}{2\cos^2\frac{V^2}{2}}(V_x^1dx - V_y^1dy),\ \ \theta_{p+3}^{p+2} = \frac{1}{2}(V_x^2dx - V_y^2dy), \tag{72}$$

$$\omega_{a'}^{a} = \lambda M_{a'}^{a}dx + \frac{1}{\lambda}N_{a'}^{a}dy$$

in (63). Then we obtain the completely integrable systems for the functions $M_{a'}^{a}, N_{a'}^{a}$:

$$M_{a',\beta_1}^{a} = 0,\ M_{p+1,V^1}^{a} = \frac{1}{2\cos\frac{V^2}{2}}M_{p+2}^{a} + \frac{\sin\frac{V^2}{2}}{2\cos^2\frac{V^2}{2}}M_{p+3}^{a},\ M_{p+1,V^2}^{a} = 0,$$

$$M^a_{p+2,V^1} = -\frac{1}{2\cos\frac{V^2}{2}}M^a_{p+1},\ M^a_{p+2,V^2} = -\frac{1}{2}M^a_{p+3},\ M^a_{p+3,V^1} = -\frac{\sin\frac{V^2}{2}}{2\cos^2\frac{V^2}{2}}M^a_{p+1},$$

$$M^a_{p+3,V^2} = \tfrac{1}{2}M^a_{p+2},\ N^a_{a',\beta_1} = 2H^a_{b\gamma_1}T^{\gamma_1}_{\beta_1}N^b_{a'},\ N^a_{p+1,V^1} = \tfrac{1}{2\cos\frac{V^2}{2}}N^a_{p+2} - \tfrac{\sin\frac{V^2}{2}}{2\cos^2\frac{V^2}{2}}N^a_{p+3},$$

$$N^a_{p+1,V^2} = 0,\ N^a_{p+2,V^1} = -\frac{1}{2\cos\frac{V^2}{2}}N^a_{p+1},\ N^a_{p+2,V^2} = \frac{1}{2}N^a_{p+3},$$

$$N^a_{p+3,V^1} = \frac{\sin\frac{V^2}{2}}{2\cos^2\frac{V^2}{2}}N^a_{p+1},\ N^a_{p+3,V^2} = -\frac{1}{2}N^a_{p+2}.$$

One can verify that integrating these systems and then substituting the found solutions and expressions (71),(72) into Eqs.(63), we obtain the Lax representation of the Euler-Lagrange system for Lagrangian (69). □

Theorem 8 *To every symmetric space* $SO(p+2)/(SO(p)\times SO(2))$ $(p\geq 3)$ *there corresponds the Lax representation of Euler's equations for Lagrangian*

$$L = [g_{\alpha_1\beta_1}(U^{\gamma_1}) + a_{\alpha_1\beta_1}(U^{\gamma_1})]U^{\alpha_1}_x U^{\beta_1}_y - 2(p-2)k^2 V_x V_y + 4(p-2)M^b_{b'}N^{b'}_b,$$

where $k = const.$

Proof. The Lax representation of these system we obtain substituting

$$\theta^a_b = H^a_{b\beta_1}T^{\beta_1}_{\gamma_1}U^{\gamma_1}_x dx,\ \theta^{p+1}_{p+2} = kV_y dy, \tag{73}$$

$$\omega^a_{a'} = \lambda M^a_{a'}dx + \frac{1}{\lambda}N^a_{a'}dy$$

in (63). Here the functions $M^a_{a'}, N^a_{a'}$ satisfy the following equations:

$$M^a_{a',\beta_1} = 0,\ M^a_{p+1,V} = kM^a_{p+2},\ M^a_{p+2,V} = -kM^a_{p+1}, \tag{74}$$

$$N^a_{a',\beta_1} = H^a_{b\gamma_1}T^{\gamma_1}_{\beta_1}N^b_{a'},\ N^a_{a',V} = 0.$$

□

Example 5.(The system associated with $SO(5)/(SO(3)\times SO(2))$.)

Choose the local coordinates and left-invariant forms on the group $SO(3)$ as in Example 1. Assume $M^1_5 = \sin U^2 \sin U^3,\ M^2_5 = -\cos U^2 \sin U^3,\ M^3_5 =$

$\cos U^3$, $M_4^a = 0$, $N_{a'}^1 = N_a^2 = 0$, $N_4^3 = l \sin kV$, $N_5^3 = l \cos kV$, $l = const$, $k = const$. Then we obtain the Lax representation of the Euler-Lagrange system for the Lagrangian

$$L = \sum_{\alpha_1=1}^{3} U_x^{\alpha_1} U_y^{\alpha_1} + k^2 V_x V_y + 2 \cos U^3 U_y^1 U_x^2 + 2l \cos U^3 \cos kV.$$

Carrying out the reductions and the Backlund transformation analogous to Example 1, we obtain the system

$$V_{xy}^1 + \frac{1}{\sin V^2} V_y^1 V_x^2 + \frac{1}{\sin V^2} V_y^2 V_x^1 = 0,$$

$$V_{xy}^2 - \frac{\sin V^2}{(1 + \cos V^2)^2} V_y^1 V_x^1 + l \sin V^2 \sin V^3 = 0, \qquad (75)$$

$$V_{xy}^3 + \frac{l}{k} \cos V^2 \sin kV^3 = 0$$

which is the Euler's system for the Lagrangian

$$L = V_x^1 V_y^1 tg^2 \frac{V^2}{2} + V_x^2 V_y^2 + k^2 V_x^3 V_y^3 + 2l \cos V^2 \cos kV^3. \qquad (76)$$

This system admits the Lax representation of the form (63), where

$$\theta_2^1 = \frac{1}{2 \cos \frac{V^2}{2}} (V_x^1 dx + V_y^1 dy), \ \theta_3^1 = -\frac{\sin \frac{V^2}{2}}{2 \cos^2 \frac{V^2}{2}} (V_x^1 dx - V_y^1 dy),$$

$$\theta_3^2 = \frac{1}{2} (V_x^2 dx - V_y^2 dy), \ \theta_5^4 = k V_x^3 dx,$$

$$\omega_{A'}^A = \lambda M_{A'}^A dx + \frac{1}{\lambda} N_{A'}^A dy,$$

$M_4^A = M_5^1 = N_4^1 = N_5^1 = 0$, $M_5^2 = -\sin \frac{V^2}{2}$, $M_5^3 = \cos \frac{V^2}{2}$, $N_4^2 = l \sin \frac{V^2}{2} \sin kV^4$, $N_5^2 = l \sin \frac{V^2}{2} \cos kV^3$, $N_4^3 = l \cos \frac{V^2}{2} \sin kV^3$, $N_5^3 = l \cos \frac{V^2}{2} \cos kV^3$.

Notice that the integrable systems with Lagrangian

$$L = V_x^1 V_y^1 tg^2 \frac{V^2}{2} + V_x^2 V_y^2 + k^2 V_x^3 V_y^3 + Q$$

were completely studied by Meshkov A.G. and Demskoi D.K. (see e.g. [1],[16]).

4 Acknowledgments

The authors are grateful to Ferapontov E.V., Stepanov N.A. and Tuzov Y.V. for helpful discussions.

References

[1] Demskoi D. K., Meshkov A. G. Zero-curvature representation for a chiral-type three-field system. Inverse Problems 19 (2003), no. 3, p. 563–571.

[2] Dijkgraaf R., Verlinde H. L., Verlinde E. P. String propagation in a black hole geometry. Nucl.Phys.B371 (1992), p. 269–314.

[3] Bilal A. Non-abelian Toda theory: a completely integrable model for strings on a black hole background. Nuclear Phys. B 422 (1994), no. 1-2, p. 258–288.

[4] Wess J., Zumino B. Consequences of anomalos ward identities. Phys.Lett.B 37(1971) P.95–97.

[5] Novikov S. P. The Hamiltonian formalism and a multivalued analogue of Morse theory. (Russian) Uspekhi Mat. Nauk 37 (1982), no. 5(227), 3–49, 248.

[6] Witten E. Nonabelian bosonization in two-dimenisions. Commun.Math.Phys.92 (1984), p. 455–472.

[7] Gawedzki K., Kupiainen A. G/H conformal field theory from gauged WZW model. Phys.Lett.B215 (1988), p.119–123.

[8] Leznov A. N., Saveliev M. V. Two-dimensional exactly and completely integrable dynamical systems. Monopoles, instantons, dual models, relativistic strings, Lund-Regge model, generalized Toda lattice, etc. Comm. Math. Phys. 89 (1983), no. 1, p. 59–75.

[9] Hollowood T. J., Miramontes J. L. Q-Han Park Massive Integrable Soliton Theories. Nucl.Phys. B445 (1995), p. 451–468.

[10] Helgason S. Differential geometry, Lie groups, and symmetric spaces. Pure and Applied Mathematics, 80, 1978, 628 pp.

[11] Zakharov V. E., Mikhailov A. V. Relativistically invariant two-dimensional models of field theory which are integrable by means of the inverse scattering problem method. (in Russian) Zh. Eksp. Teor. Fiz.74 (1978), p. 1953–1973.

[12] Balandin A. V. A system of differential equations that admits a zero curvature representation. (in Russian) Uspekhi Mat. Nauk 45 (1990), no. 6(276), p. 125–126; translation in Russian Math. Surveys 45 (1990), no. 6, p. 141–142.

[13] Balandin A.V., Pakhareva O.N., Potemin G.V. Lax representation of the chiral-type field equations. Phys. Lett. A. 2001. V. 283, N. 3-4. P. 168—176.

[14] Getmanov B.S. An integrable two-dimensional Lorentz-invariant nonlinear model of a complex scalar field (complex sine-Gordon-II). (in Russian) Teoret. Mat. Fiz. 1981. V. 48. N. 1, p.13-23.

[15] Witten E. On string theory and black holes. Phys.Rev.D44 (1991), p. 314–324

[16] Demskoi D.K., Meshkov A.G. The Lax representation for a triplet of scalar fields. (in Russian) Teoret. Mat. Fiz. 2003. V. 134, N. 3, p. 401.